

ONLINE FIRST

Sustained Decreases in Risk Exposure and Youth Problem Behaviors After Installation of the Communities That Care Prevention System in a Randomized Trial



Scan for Author
Audio Interview

J. David Hawkins, PhD; Sabrina Oesterle, PhD; Eric C. Brown, PhD; Kathryn C. Monahan, PhD; Robert D. Abbott, PhD; Michael W. Arthur, PhD; Richard F. Catalano, PhD

Objective: To test whether the Communities That Care (CTC) prevention system reduced levels of risk and adolescent problem behaviors community-wide 6 years after installation of CTC and 1 year after study-provided resources ended.

Design: A community randomized trial.

Setting: Twenty-four small towns in 7 states, matched within state, randomly assigned to control or intervention condition in 2003.

Participants: A panel of 4407 fifth-grade students was surveyed annually through 10th grade from 2004 to 2009.

Intervention: A coalition of community stakeholders received training and technical assistance to install CTC, used epidemiologic data to identify elevated risk factors and depressed protective factors in the community, and implemented programs to address their community's elevated risks from a menu of tested and effective programs for youths aged 10 to 14 years, their families, and schools.

Outcome Measures: Levels of risk and incidence and prevalence of tobacco, alcohol, and other drug use; delinquency; and violent behavior by grade 10.

Results: Mean levels of targeted risks increased less rapidly between grades 5 and 10 in CTC than in control communities and were significantly lower in CTC than control communities in grade 10. The incidence of delinquent behavior, alcohol use, and cigarette use and the prevalence of current cigarette use and past-year delinquent and violent behavior were significantly lower in CTC than in control communities in grade 10.

Conclusions: Using the CTC system can produce enduring reductions in community-wide levels of risk factors and problem behaviors among adolescents beyond the years of supported implementation, potentially contributing to long-term public health benefits.

Trial Registration: clinicaltrials.gov Identifier: NCT01088542

Arch Pediatr Adolesc Med. 2012;166(2):141-148.
Published online October 3, 2011.
doi:10.1001/archpediatrics.2011.183

Author Affiliations: Social Development Research Group, School of Social Work (Drs Hawkins, Oesterle, Brown, Monahan, Arthur, and Catalano), and Department of Educational Psychology (Dr Abbott), University of Washington, Seattle. Dr Monahan is now with the Department of Psychology, University of Pittsburgh, Pittsburgh, Pennsylvania.

SMOKING, DRINKING, AND VIOLENCE are among the leading preventable causes of death in the United States and have large costs to society.¹⁻⁵ The prevention of delinquency, violence, and the use of alcohol, tobacco, and other drugs during early adolescence are important public health priorities because involvement in these behaviors often begins in childhood and adolescence,⁶⁻⁸ and early initiation of these behaviors is associated with greater risk for later serious health and behavior problems.⁹⁻¹⁵

Advances in prevention science over the past 2 decades have identified common risk factors for adolescent substance use and delinquent and violent behavior and have pro-

duced a growing list of tested and effective preventive interventions,^{16,17} yet these advances are not systematically used by local decision makers.¹⁸⁻²⁰ Widespread installation of tested and effective preventive interventions could have significant benefits to public health.^{6,21,22} Systems for increasing use of tested and effective prevention programs in communities are needed. Communities That Care (CTC)²³ is a system for guiding communities to choose, install, and monitor tested and effective preventive interventions to address elevated risks and suppressed protective factors affecting youth.

Communities That Care mobilizes community stakeholders to collaborate on the development and implementation of a sci-

ence-based community prevention system. The CTC system is expected to produce community-wide changes in prevention system functioning, including increased adoption of a science-based approach to prevention and increased use of tested, effective preventive interventions that address risk and protective factors prioritized by the community. These changes in prevention system functioning are expected to produce community-wide changes in youths' exposure to targeted risk and protective factors, which, in turn, result in decreases in adolescent substance use and delinquent behaviors. The CTC theory of change suggests that it takes 2 to 5 years of implementing tested, effective programs for community-level impact on risk and protective factors and 4 to 10 years for community-level impact on adolescent substance use, delinquency, and violence to be observed.²⁴

Results from the first randomized trial of CTC^{25,26} found support for the CTC theory of change²⁷⁻³¹ and demonstrated that 4 years after the initial implementation of CTC the incidences of delinquent behavior, alcohol use, cigarette use, and smokeless tobacco use were significantly lower in CTC than in control communities by the end of eighth grade in a panel of students followed up from fifth grade. In addition, grade 8 prevalence rates of alcohol and smokeless tobacco use in the past 30 days, binge drinking in the past 2 weeks, and the number of different delinquent behaviors committed in the past year were significantly lower in CTC than in control communities.³²

Communities That Care has been placed in the public domain by the Substance Abuse and Mental Health Services Administration and is available at www.communitiesthatcare.net. Given the potential for widespread dissemination of the CTC system, it is important to understand the degree to which installation of CTC during the initial 5-year efficacy trial led to sustained differences in outcomes between CTC and control communities beyond the end of study-provided resources that supported the installation of CTC.³³⁻³⁶ The present study tested CTC's effects on youth problem behaviors 1 year after study-provided resources ended, ie, 6 years after initial implementation of CTC and 4.67 years after CTC communities began implementing prevention programs selected through the CTC process.

METHODS

The Community Youth Development Study (CYDS)²⁶ is the first community randomized trial of CTC. Twenty-four communities in the states of Colorado, Illinois, Kansas, Maine, Oregon, Utah, and Washington were matched in pairs within state on population size, racial and ethnic diversity, economic indicators, and crime rates. One community from within each matched pair was assigned randomly by a coin toss to either the intervention (CTC) or control condition.²⁶ The CYDS communities are small- to moderate-sized incorporated towns with their own governmental, educational, and law enforcement structures ranging in population from 1500 to 50 000 residents.

Beginning in the summer of 2003, intervention communities received 6 CTC training sessions delivered over the course of 6 to 12 months by certified CTC trainers. The CTC coalition members were trained to use data from surveys of public

school students in the community to prioritize risk factors to be targeted by tested and effective preventive actions in the community.^{37,38} For the CYDS, CTC communities were asked to focus their prevention plans on programs for youths aged 10 to 14 years (grades 5-9) and their families and schools so that possible effects on drug use and delinquency could be observed within the initial 5-year grant period. Starting with the 2004-2005 school year and annually thereafter, community coalitions implemented between 1 and 5 preventive programs to address their prioritized risk factors. The CYDS implementation staff provided technical assistance throughout the 5-year efficacy trial (2003-2008) via weekly telephone calls, e-mails, and site visits to CTC communities at least once per year. Technical assistance ended after the fifth year of the study. Control communities received data from the CTC Youth Surveys administered in their schools every 2 years but no other training or technical assistance from the study.

SAMPLE AND DATA COLLECTION

Data were drawn from a longitudinal panel of public school students in the 24 CYDS communities followed up annually from grade 5 through grade 10 (N=4407; 50% male; 67% white; 20% Hispanic).²⁵ All students in fifth-grade classrooms during the 2003-2004 school year in these schools were eligible for participation in the CYDS. The first wave of data collection (fifth grade, 2004) was a preintervention baseline assessment. Recruitment continued in wave 2 (grade 6) to increase the overall participation rate. This strategy resulted in a total of 4420 students whose parents consented to their participation in the study (76.4% of the eligible population). The sixth annual wave of data was collected in the spring of 2009 when students were in grade 10, approximately 4.67 years after the prevention programs chosen by CTC communities were first implemented. At this point, CTC communities had not received any technical assistance for 1 year.

The active longitudinal panel consists of 4407 students who completed a wave 1 or wave 2 survey. Students in the longitudinal panel who remained in the intervention or control communities for at least 1 semester were tracked and surveyed annually, even if they left the community.²⁵ Ninety-four percent of students in the longitudinal panel completed the survey in wave 6 (10th grade). **Figure 1** shows the flow of communities and students through the study.

Students completed the Youth Development Survey,³⁹ a self-administered paper-and-pencil questionnaire designed to be completed in a class period. Identification numbers but no names or other identifying information were included on the surveys. The University of Washington Human Subjects Review Committee approved this protocol.

MEASURES

Targeted Risk Factors

Based on anonymous baseline surveys of all assenting sixth- and eighth-grade students in each CTC community, CTC coalitions prioritized between 2 and 5 risk factors that were elevated in their community.^{31,40} An average targeted risk factor score was calculated by standardizing and averaging the community-specific set of targeted risk factors in CTC communities. Because control communities did not prioritize and target risk and protective factors using the CTC process, each control community's average risk factor score was calculated using the set of targeted risk factors identified in its matched CTC community.




Figure 1. Flow of communities and participants in the randomized trial.

Substance Use

Students self-reported the incidence of first use of alcohol, cigarettes, smokeless tobacco, marijuana, inhalants, and prescription drugs not prescribed by a doctor between grades 5 and 10. The prevalence of binge drinking during the last 2 weeks and use of alcohol, cigarettes, smokeless tobacco, marijuana, and inhalants in the last month were measured in grades 5 and 10. In grade 10, the prevalence rates of illicit use of prescription drugs and other illicit drugs (ie, psychedelics, 3,4-methylenedioxymethamphetamine [MDMA or Ecstasy], stimulants, and cocaine or crack) in the past month also were assessed.

Delinquent and Violent Behavior

The incidence of delinquent behavior was operationalized as the first self-reported occurrence of any of 4 delinquent acts (stealing, property damage, shoplifting, or attacking someone with intent to harm) between grades 5 and 10. Questions about more serious delinquent behavior (carrying a gun to school, beating someone up, stealing a vehicle, selling drugs, or being arrested) were added to the survey over time as developmentally appropriate. A measure of the variety of delinquent acts committed in the past year was calculated in the fifth grade (range, 0-4) and 10th grade (range, 0-9). A subset of the delinquency items was used to create a measure of violent be-




Figure 2. Unadjusted average targeted risk factor score by intervention condition. CTC indicates Communities That Care.

havior in the fifth grade (attacking someone with intent to harm; range, 0-1) and 10th grade (attacking someone with intent to harm, carrying a gun to school, beating somebody up; range, 0-3). There were no statistically significant differences in levels of average targeted risk factors or the prevalence or mean number of delinquent and violent acts by intervention status at baseline.^{25,31}

Student and Community Characteristics

Characteristics of students and communities were used as covariates in all analyses. Student-level covariates included age at time of the grade 6 survey, race, Hispanic ethnicity, parental education, attendance at religious services during grade 5 (1=never to 4=about once a week or more), and rebelliousness in grade 5 (mean of 3 items; $\alpha = .69$). Community-level covariates included total population of students in the community and the percentage of students eligible for free or reduced-price school lunch.

ANALYSIS SAMPLE AND MISSING DATA PROCEDURES

Of the 4407 students in the longitudinal panel, 26.5% were recruited at wave 2 (in grade 6) and thus did not complete a questionnaire in wave 1 (grade 5). Overall, 93.5% of the students participated in at least 5 of the 6 waves of data collection. There was no systematic bias from differential accretion or differential attrition in control and intervention conditions.²⁵ Students were excluded from the analysis if they reported being honest “some of the time” or less on the survey, reported having used a fictitious drug included in the survey as a validity screen, or reported that they had used 2 of 3 drugs (marijuana, inhalants, or other drugs) on 40 occasions or more during the past month.³² Based on these validity criteria, 66 students were excluded from analyses in grade 10, resulting in valid data from 4069 students in 10th grade.

Item nonresponse was small (<1%). Missing data were imputed using multiple imputations.⁴¹ Using NORM version 2.03,⁴² 40 separate data sets including data from all 6 waves were imputed separately by intervention condition.⁴³ Analyses were conducted within each imputed data set and results were averaged across all data sets using the Rubin rules.⁴⁴

ANALYSES

Differential change in levels of average targeted risk by intervention condition from grade 5 to grade 10 was assessed using 3-level latent growth models^{45,46} to account for nesting of time within students and students within communities. Characteristics of students and communities, and 11 dummy variables indicating the matching of communities into pairs, were included as model covariates. Analyses were conducted using HLM version 6.08.⁴⁷

Multilevel discrete time survival analysis^{48,49} was used to examine the effect of CTC on the incidence of drug use and delinquency between grades 5 and 10. The risk of initiating drug use and delinquent behavior was assessed for students who had not yet initiated these behaviors before the grade 5 survey. To test if the effect of the intervention on incidence was constant across time, we included interaction effects between intervention condition and time. All analyses included student and community covariates and were analyzed using MLwiN version 2.02.⁵⁰

The effect of CTC on grade 10 prevalence rates of substance use, delinquency, and violence was assessed using the generalized linear mixed model^{51,52} and analyzed using HLM version 6.08⁴⁷ using a logit link transformation for the Bernoulli distributed drug use and delinquency outcomes and using a log link function for the Poisson distributed measures of the variety of delinquent and violent acts. In addition to student and community characteristics, the respective preintervention drug use, delinquency, or violence measure was included as a baseline covariate.

Intervention effects were estimated as the mean differences in adjusted community-level slopes, means, and rates between intervention and control communities as tested against the average variation among intervention condition-specific adjusted community-level slopes, means, and rates. The *df* for intervention effects were equal to the number of community-matched pairs (12) minus the number of community-level covariates, minus 1. Statistically significant effects were assessed at a .05 type I error rate (2-tailed). To be consistent with prior reports of findings from this study,³² we report adjusted odds ratios (AORs), which were almost identical to corrected risk ratios.⁵³

RESULTS

TARGETED RISK

Unadjusted mean levels of the average targeted risk factor score from grade 5 to grade 10 by intervention condition are shown in **Figure 2**. The latent growth model indicated that the increase in targeted risk between grades 5 and 10 was significantly smaller in CTC than in control communities ($t_{11} = -2.43$; $P < .05$). The standardized effect size corresponding to this intervention effect was $\delta = -1.76$ SDs.⁵⁴ Mean levels of targeted risk were equivalent in CTC and control communities at baseline in grade 5 ($t_0 = 1.04$; $P > .05$) but were significantly lower in CTC than in control communities at grade 10 ($t_9 = -2.54$; $P < .05$; Cohen $d = -0.12$).

INCIDENCE OF DRUG USE AND DELINQUENCY

Results from the multilevel discrete time survival analysis indicated a significant effect of CTC in reducing the incidence of the use of alcohol and cigarettes by 10th grade




Figure 3. Unadjusted cumulative initiation of substance use by intervention condition. CTC indicates Communities That Care.

among students who had not yet initiated use by fifth grade. **Figure 3** shows the unadjusted cumulative initiation rates for each drug by intervention condition. The AOR for the effect of CTC on alcohol use incidence in 10th grade was 0.62 (95% confidence interval [CI], 0.41-0.94; $t_{11} = -2.53$; $P = .03$), indicating that students in CTC communities had 38% lower odds of initiating the use of alcohol in grade 10 than students in control communities. The odds of initiating cigarette use were significantly different between CTC and control communities in 10th grade, with an AOR of 0.54 (95% CI, 0.36-0.81; $t_{11} = -3.36$; $P = .006$), suggesting that students in CTC communities had almost half the odds of beginning to smoke in 10th grade as students in control communities. Significant differences by intervention status in the incidence of smokeless tobacco, marijuana, inhalant, or prescription drug use were not observed by the spring of 10th grade (Figure 3).

Figure 4 shows unadjusted cumulative initiation rates of delinquent behavior by intervention condition. Re-




Figure 4. Unadjusted cumulative initiation of delinquent behavior by intervention condition. CTC indicates Communities That Care.

sults indicated a significant effect of CTC on the incidence of delinquent behavior between grades 6 and 10. The effect of CTC on the incidence of delinquent behavior did not vary across time. The AOR for initiating any

Table. Unadjusted Prevalence Rates of Current Drug Use and Delinquency in Grade 10 and AORs Comparing Control and CTC Communities

	%		AOR (95% CI) ^a
	Control Communities	CTC Communities	
Drug use			
Last 30 d			
Alcohol	29.5	28.7	1.10 (0.82-1.47)
Cigarettes	16.3	13.4	0.79 (0.64-0.99) ^b
Smokeless tobacco	7.5	6.3	0.85 (0.63-1.15)
Inhalants	2.0	3.1	1.50 (0.88-2.58)
Marijuana	15.0	14.5	0.99 (0.66-1.49)
Prescription drugs	4.8	5.6	1.15 (0.81-1.65)
Other illicit drugs	5.5	6.5	1.25 (0.90-1.73)
Last 2 wk			
Binge drinking	13.6	11.0	0.89 (0.67-1.19)
Delinquent behavior in last year			
No. of delinquent behaviors, mean	1.1	0.9	0.89 (0.77-1.03)
Any delinquency	42.6	36.1	0.83 (0.69-0.99) ^b
Violent behavior in last year			
No. of violent behaviors, mean	0.3	0.2	0.84 (0.66-1.05)
Any violence	17.7	13.2	0.75 (0.58-0.97) ^b

Abbreviations: AOR, adjusted odds ratio; CI, confidence interval; CTC, Communities That Care.

^aOdds ratios are adjusted for grade 5 prevalence, student age, sex, race, ethnicity, parental education, grade 5 religious attendance, grade 5 rebelliousness, student population of the community, and percentage of students in the community receiving free or reduced-price school lunch. Alcohol use in the last 30 days in grade 5 was used to adjust analyses of 10th-grade marijuana, prescription drug, and other illicit drug use because of very low baseline use of these drugs.

^b $P < .05$

delinquent act between grades 6 and 10 was 0.79 (95% CI, 0.64-0.98, $t_0 = -2.52$; $P = .03$), indicating that the odds of initiating delinquent behavior by 10th grade were 21% lower for students in CTC communities than students in control communities.

PREVALENCE OF DRUG USE

Unadjusted prevalence rates and AORs of current drug use in the 10th grade in CTC and control communities are presented in the **Table**. Results from the multivariate analyses indicated significantly lower prevalence rates in the 10th grade in CTC communities compared with control communities for past-month cigarette use ($t_0 = -2.38$; $P = .04$; AOR=0.79). The odds of smoking cigarettes in the past month were 21% lower for students in CTC communities than students in control communities. Tenth graders in control and CTC communities did not differ significantly in rates of binge drinking in the past 2 weeks or in past-month alcohol, smokeless tobacco, marijuana, inhalant, prescription, or other illicit drug use.

PREVALENCE AND VARIETY OF DELINQUENT AND VIOLENT BEHAVIORS

The Table shows that 10th-grade students in CTC communities had 17% lower odds of reporting any delinquent behavior in the past year ($t_0 = -2.33$; $P = .04$; AOR=0.83) and 25% lower odds of reporting any violent behavior in the past year ($t_0 = -2.51$; $P = .03$; AOR=0.75) compared with students in control communities. However, the variety of different delinquent or violent acts in which students engaged was not significantly lower among students in CTC communities

compared with students in the control communities (delinquency: $t_0 = -1.86$; $P = .10$; AOR=0.89; violence: $t_0 = -1.75$; $P = .10$; AOR=0.84).

COMMENT

The CTC prevention system previously has been found to decrease risk factors and youths' involvement in adolescent substance use and delinquent behavior during a period of study-supported intervention.³² We have conducted analyses of the ratio of economic benefits of these outcomes to the costs of CTC implementation, which indicate that the benefits of CTC exceed its costs; these results are reported in a separate article. The present study found that community-wide beneficial effects of the CTC prevention system on the incidence of adolescent delinquency and tobacco and alcohol use and the prevalence of delinquent behavior, violence, and tobacco use were observed 1 year beyond supported implementation of CTC. These are the most prevalent adolescent problem behaviors of those measured in this study and have the greatest costs to society. The use of CTC did not eliminate these adolescent problem behaviors in participating communities, but it reduced them significantly. It will be important to compare the economic value of these sustained benefits with the costs of implementing CTC. By the end of grade 10, the use of CTC did not have significant effects on the variety of different delinquent or violent acts and less prevalent adolescent problem behaviors including the use of marijuana or other illegal drugs. It is possible that effects on the use of these drugs may be observed later as their use becomes more prevalent in the adolescent population.

An important characteristic of the CTC system is that it guides coalitions of community stakeholders to assess the prevalence of exposure to empirically identified risk and protective factors among young people in the community and to choose tested and effective preventive interventions to reduce those risk factors that are most prevalent in each community. Each community in this study selected a different set of preventive interventions to address that community's unique profile of risk and protection. This study found that using the CTC system to diagnose and address each community's prevention needs in this way had community-wide effects in reducing alcohol and tobacco use and delinquent and violent behavior.

These findings from the randomized trial, coupled with quasiexperimental results from using CTC in Pennsylvania,^{55,56} indicate that the use of CTC can contribute to long-term community-wide improvements in public health. We have developed a system for training new CTC trainers and technical assistance providers that can be offered to state agencies and other organizations that seek to build capacity to provide CTC to communities. This system ensures that new trainers have the opportunity to provide CTC training to communities under the supervision of certified CTC trainers and to become certified CTC trainers through this process. The availability of this system increases the likelihood that CTC can be disseminated widely and, if disseminated, will have widespread public health benefits.

Accepted for Publication: July 13, 2011.

Published Online: October 3, 2011. doi:10.1001/archpediatrics.2011.183

Correspondence: J. David Hawkins, PhD, Social Development Research Group, University of Washington, 9725 3rd Ave NE, Ste 401, Seattle, WA 98115 (jdh@uw.edu).

Author Contributions: Dr Hawkins had full access to all of the data in the study and takes responsibility for the integrity of the data and the accuracy of the data analysis. *Study concept and design:* Hawkins, Oesterle, Brown, Monahan, Abbott, Arthur, and Catalano. *Acquisition of data:* Hawkins, Arthur, and Catalano. *Analysis and interpretation of data:* Hawkins, Oesterle, Brown, Monahan, Abbott, Arthur, and Catalano. *Drafting of the manuscript:* Hawkins, Oesterle, Brown, and Monahan. *Critical revision of the manuscript for important intellectual content:* Hawkins, Oesterle, Brown, Monahan, Abbott, Arthur, and Catalano. *Statistical analysis:* Oesterle, Brown, Monahan, and Abbott. *Obtained funding:* Hawkins, Arthur, and Catalano. *Administrative, technical, and material support:* Hawkins and Oesterle. *Study supervision:* Hawkins, Arthur, and Catalano.

Financial Disclosure: Dr Catalano is a board member of Channing Bete Company, distributor of Supporting School Success and Guiding Good Choices. These programs were used in some communities in the study that produced the data set used in this article.

Funding/Support: This work was supported by research grant R01 DA015183-03 from the National Institute on Drug Abuse, with cofunding from the National Cancer Institute, the National Institute of Child Health and Human Development, the National Institute of Mental Health, the Center for Substance Abuse Prevention,

and the National Institute on Alcohol Abuse and Alcoholism.

Role of the Sponsor: The funding organizations had no role in the design and conduct of the study; collection, analysis, or preparation of data; or preparation, review, or approval of the manuscript.

Disclaimer: The content of this article is solely the responsibility of the authors and does not necessarily represent the official views of the funding agencies.

Additional Contributions: David M. Murray, PhD, Ohio State University, provided paid statistical consultation on this project, but the authors are responsible for all analyses and results. We acknowledge the contributions of the communities participating in the Community Youth Development Study and the collaborating state offices of drug abuse prevention in Colorado, Illinois, Kansas, Maine, Oregon, Utah, and Washington.

REFERENCES

- Mokdad AH, Marks JS, Stroup DF, Gerberding JL. Actual causes of death in the United States, 2000. *JAMA*. 2004;291(10):1238-1245.
- Woolf SH. The big answer: rediscovering prevention at a time of crisis in health care. *Harv Health Policy Rev*. 2006;7(2):5-20.
- Specter A. Making youth violence prevention a national priority. *Am J Prev Med*. 2008;34(3)(suppl):S3-S4.
- US Department of Health and Human Services. *Youth Violence: A Report of the Surgeon General*. Rockville, MD: US Dept of Health and Human Services, Substance Abuse and Mental Health Services Administration, Center for Mental Health Services, National Institutes of Health, National Institute of Mental Health; 2001.
- McCullister KE, French MT, Fang H. The cost of crime to society: new crime-specific estimates for policy and program evaluation. *Drug Alcohol Depend*. 2010;108(1-2):98-109.
- Woolf SH. Potential health and economic consequences of misplaced priorities. *JAMA*. 2007;297(5):523-526.
- Centers for Disease Control and Prevention. The health consequences of smoking: a report of the Surgeon General. <http://www.surgeongeneral.gov/library/smokingconsequences>. Published May 24, 2004. Accessed July 8, 2010.
- Substance Abuse and Mental Health Services Administration—US Department of Health and Human Services. Report to Congress: a comprehensive plan for preventing and reducing underage drinking. <http://store.samhsa.gov/shin/content//SMA11-4645/SMA11-4645.pdf>. Accessed August 17, 2011.
- Grant JD, Scherrer JF, Lynskey MT, et al. Adolescent alcohol use is a risk factor for adult alcohol and drug dependence: evidence from a twin design. *Psychol Med*. 2006;36(1):109-118.
- Robins LN, Przybeck TR. Age of onset of drug use as a factor in drug and other disorders. In: Jones CL, Battjes RJ, eds. *Etiology of Drug Abuse: Implications for Prevention*. Rockville, MD: National Institute on Drug Abuse; 1985:178-192. NIDA Research Monograph 56.
- Hingson RW, Heeren T, Winter MR. Age at drinking onset and alcohol dependence: age at onset, duration, and severity. *Arch Pediatr Adolesc Med*. 2006;160(7):739-746.
- US Department of Health and Human Services. The Surgeon General's call to action to prevent and reduce underage drinking. <http://www.surgeongeneral.gov/topics/underagedrinking>. Accessed July 8, 2010.
- Agrawal A, Sartor CE, Lynskey MT, et al. Evidence for an interaction between age at first drink and genetic influences on DSM-IV alcohol dependence symptoms. *Alcohol Clin Exp Res*. 2009;33(12):2047-2056.
- Sampson RJ, Laub JH. Life-course desisters? trajectories of crime among delinquent boys followed to age 70. *Criminology*. 2003;41(3):555-592. doi:10.1111/j.1745-9125.2003.tb00997.x.
- Farrington DP. Developmental and life-course criminology: key theoretical and empirical issues—the 2002 Sutherland Award Address. *Criminology*. 2003;41(2):221-255. doi:10.1111/j.1745-9125.2003.tb00987.x.
- Mihalic S, Fagan A, Irwin K, Ballard D, Elliott D. *Blueprints for Violence Prevention*. Boulder: National Youth Violence Prevention Resource Center, University of Colorado; 2004.
- O'Connell ME, Boat T, Warner KE. *Preventing Mental, Emotional, and Behavioral Disorders Among Young People: Progress and Possibilities*. Washington, DC: National Academies Press; 2009.

18. Hallfors D, Godette D. Will the 'principles of effectiveness' improve prevention practice? early findings from a diffusion study. *Health Educ Res.* 2002;17(4):461-470.
19. Gottfredson DC, Gottfredson GD. Quality of school-based prevention programs: results from a national survey. *J Res Crime Delinq.* 2002;39(1):3-35. doi:10.1177/002242780203900101.
20. Ringwalt CL, Ennett S, Vincus A, Thorne J, Rohrbach LA, Simons-Rudolph A. The prevalence of effective substance use prevention curricula in U.S. middle schools. *Prev Sci.* 2002;3(4):257-265.
21. Woolf SH. The power of prevention and what it requires. *JAMA.* 2008;299(20):2437-2439.
22. Woolf SH. The meaning of translational research and why it matters. *JAMA.* 2008;299(2):211-213.
23. Hawkins JD, Catalano RF. *Investing in Your Community's Youth: An Introduction to the Communities That Care System.* South Deerfield, MA: Channing Bete; 2002.
24. Hawkins JD, Catalano RF. *Communities That Care Community Board Orientation: Participant's Guide.* www.communitiesthatcare.net. Accessed July 12, 2010.
25. Brown EC, Graham JW, Hawkins JD, et al. Design and analysis of the Community Youth Development Study longitudinal cohort sample. *Eval Rev.* 2009;33(4):311-334.
26. Hawkins JD, Catalano RF, Arthur MW, et al. Testing Communities That Care: the rationale, design and behavioral baseline equivalence of the Community Youth Development Study. *Prev Sci.* 2008;9(3):178-190.
27. Quinby RK, Fagan AA, Hanson K, Brooke-Weiss B, Arthur MW, Hawkins JD. Installing the Communities That Care prevention system: implementation progress and fidelity in a randomized controlled trial. *J Comp Psychol.* 2008;36(3):313-332. doi:10.1002/jcop.20194.
28. Fagan AA, Hanson K, Hawkins JD, Arthur MW. Translational research in action: implementation of the Communities That Care prevention system in 12 communities. *J Comp Psychol.* 2009;37(7):809-829. doi:10.1002/jcop.20332.
29. Brown EC, Hawkins JD, Arthur MW, Briney JS, Abbott RD. Effects of Communities That Care on prevention services systems: findings from the Community Youth Development Study at 1.5 years. *Prev Sci.* 2007;8(3):180-191.
30. Fagan AA, Hanson K, Hawkins JD, Arthur MW. Implementing effective community-based prevention programs in the Community Youth Development Study. *Youth Violence Juv Justice.* 2008;6(3):256-278. doi:10.1177/1541204008315937.
31. Hawkins JD, Brown EC, Oesterle S, Arthur MW, Abbott RD, Catalano RF. Early effects of Communities That Care on targeted risks and initiation of delinquent behavior and substance use. *J Adolesc Health.* 2008;43(1):15-22.
32. Hawkins JD, Oesterle S, Brown EC, et al. Results of a type 2 translational research trial to prevent adolescent drug use and delinquency: a test of Communities That Care. *Arch Pediatr Adolesc Med.* 2009;163(9):789-798.
33. Shediach-Rizkallah MC, Bone LR. Planning for the sustainability of community-based health programs: conceptual frameworks and future directions for research, practice and policy. *Health Educ Res.* 1998;13(1):87-108.
34. Swisher JD, Clayton R. Sustainability of prevention. *Addict Behav.* 2000;25(6):965-973.
35. Gomez BJ, Greenberg MT, Feinberg ME. Sustainability of community coalitions: an evaluation of Communities That Care. *Prev Sci.* 2005;6(3):199-202.
36. Steckler A, Goodman RM. How to institutionalize health promotion programs. *Am J Health Promot.* 1989;3(4):34-44.
37. Arthur MW, Hawkins JD, Pollard JA, Catalano RF, Baglioni AJ Jr. Measuring risk and protective factors for substance use, delinquency, and other adolescent problem behaviors: the Communities That Care Youth Survey. *Eval Rev.* 2002;26(6):575-601.
38. Glaser RR, Van Horn ML, Arthur MW, Hawkins JD, Catalano RF. Measurement properties of the Communities That Care® Youth Survey across demographic groups. *J Quant Criminol.* 2005;21(1):73-102. doi:10.1007/s10940-004-1788-1.
39. Social Development Research Group. *Community Youth Development Study, Youth Development Survey (Grades 5-10).* Seattle: Social Development Research Group, School of Social Work, University of Washington; 2005-2009.
40. Oesterle S, Hawkins JD, Fagan AA, Abbott RD, Catalano RF. Testing the universality of the effects of the Communities That Care prevention system for preventing adolescent drug use and delinquency. *Prev Sci.* 2010;11(4):411-423.
41. Schafer JL, Graham JW. Missing data: our view of the state of the art. *Psychol Methods.* 2002;7(2):147-177.
42. Schafer JL. *NORM for Windows 95/98/NT Version 2.03.* University Park, PA: Center for the Study and Prevention Through Innovative Methodology at Pennsylvania State University; 2000.
43. Graham JW, Taylor BJ, Olchowski AE, Cumsille PE. Planned missing data designs in psychological research. *Psychol Methods.* 2006;11(4):323-343.
44. Rubin DB. *Multiple Imputation for Nonresponse in Surveys.* New York, NY: Wiley; 1987.
45. Laird NM, Ware JH. Random-effects models for longitudinal data. *Biometrics.* 1982;38(4):963-974.
46. Raudenbush SW. Toward a coherent framework for comparing trajectories of individual change. In: Collins LM, Sayer AG, eds. *New Methods for the Analysis of Change.* Washington, DC: American Psychological Association; 2001:35-64.
47. Raudenbush SW, Bryk AS. *Hierarchical Linear Models: Applications and Data Analysis Methods.* 2nd ed. Newbury Park, CA: Sage; 2002.
48. Barber JS, Murphy S, Axinn WG, Maples J. Discrete-time multilevel hazard analysis. *Sociol Methodol.* 2000;30(1):201-235. doi:10.1111/0081-1750.00079.
49. Reardon SF, Brennan RT, Buka SL. Estimating multi-level discrete-time hazard models using cross-sectional data: neighborhood effects on the onset of adolescent cigarette use. *Multivariate Behav Res.* 2002;37(3):297-330. doi:10.1207/S15327906MBR3703_1.
50. Rasbash J, Browne W, Healy M, Cameron B, Charlton C. *MLwiN Version 2.02.* Bristol, England: Multilevel Models Project, Institute of Education, University of Bristol; 2004.
51. Breslow N, Clayton DG. Approximate inference in generalized linear mixed models. *J Am Stat Assoc.* 1993;88(421):9-25. doi:10.2307/2290687.
52. Liang KY, Zeger SL. Longitudinal data analysis using generalized linear models. *Biometrika.* 1986;73(1):13-22. doi:10.2307/2336267.
53. Zhang J, Yu KF. What's the relative risk? a method of correcting the odds ratio in cohort studies of common outcomes. *JAMA.* 1998;280(19):1690-1691.
54. Raudenbush SW, Xiao-Feng L. Effects of study duration, frequency of observation, and sample size on power in studies of group differences in polynomial change. *Psychol Methods.* 2001;6(4):387-401.
55. Feinberg ME, Jones D, Greenberg MT, Osgood DW, Bontempo D. Effects of the Communities That Care model in Pennsylvania on change in adolescent risk and problem behaviors. *Prev Sci.* 2010;11(2):163-171.
56. Feinberg ME, Greenberg MT, Osgood DW, Sartorius J, Bontempo D. Effects of the Communities That Care model in Pennsylvania on youth risk and problem behaviors. *Prev Sci.* 2007;8(4):261-270.

Copyright of Archives of Pediatrics & Adolescent Medicine is the property of American Medical Association and its content may not be copied or emailed to multiple sites or posted to a listserv without the copyright holder's express written permission. However, users may print, download, or email articles for individual use.